arXiv:1510.02423v1 [math.RT] 8 Oct 2015

Representations of the discrete Heisenberg group on distribution spaces of two-dimensional local fields

D. V. Osipov, A. N. Parshin *

To Vladimir Petrovich Platonov on his 75th birthday

Abstract

We study a natural action of the Heisenberg group of integer unipotent matrices of the third order on distribution space of a two-dimensional local field for a flag on a two-dimensional scheme.

1 Introduction

Let X be an algebraic curve over a field k. Then one can attach a discrete valuation ring $\mathcal{O}_x$ and a local field $K_x = \mathrm{Frac}(\mathcal{O}_x)$ to a closed point $x \in X$. Thus the valuation group $\Gamma_x = K_x^*/\mathcal{O}_x^* = \mathbb{Z}$ will be the simplest discrete Abelian group canonically attached to this point of the curve. Starting from this local group one can define the global group $\mathrm{Div}(X) = \oplus_{x\in X}\Gamma_x$.

In arithmetics, when $k = \mathbb{F}_q$, the representation theory of these groups plays an important role in the study of arithmetical invariants of the curve X such as number of classes or zeta-function. These groups act on the functional spaces of the field K_x and of the adelic group $\mathbb{A}_X$. All irreducible representations are of dimension 1 (i.e. characters with values in $\mathbb{C}^*$), and the functional spaces which are infinite-dimensional can be presented as certain integrals (but not direct sums) of these characters. This construction is very useful if one wants to get an analytical continuation and functional equation of L-functions of the curve X (see [14, 15, 13]).

The goal of this work is to extend this circle of questions to algebraic surfaces. We will restrict ourselves by the local groups only. As it was shown in [12] and in the previous publications, the local fields will now be the two-dimensional local fields, and the corresponding discrete group is the simplest non-Abelian nilpotent group of class 2: the Heisenberg group $\mathrm{Heis}(3,\mathbb{Z})$ of unipotent matrices of order 3 with integer entries. This group has a very non-trivial representation theory (see [11, 12]).

*Both authors are partially supported by Russian Foundation for Basic Research (grants no. 13-01-12420 ofi_m2 and no. 14-01-00178a) and by the grant of supporting of Leading Scientific Schools NSh-2998.2014.1.

The classical representation theory of unitary representations of localy compact groups in a Hilbert space does not work smoothly at this case. The Heisenberg group is not a group of type I in the von Neumann classification. Thus, the Heisenberg group has a bad topology in its unitary dual space, has, generally speaking, no characters for irreducible representations, and the decompositions of representations into integrals of the irreducible ones can be non-unique, see [4]. Note that Abelian groups, compact topological groups and semi-simple Lie groups are the groups of type I. In our case, if we change the category of representation spaces and consider instead of the Hilbert spaces the vector spaces of countable dimension and without any topology, then the situation will be much better. This approach is modeled onto the theory of smooth representations of reductive algebraic groups over non-Archimedean local fields. They are so called l-groups, and the discrete groups are such groups. Then the basic notions of the smooth representation theory (see, for example, [3]) can be applied in this case. The new approach get for smooth non-unitary representations a moduli space which is a complex manifold. The characters can be defined as modular forms on this manifold, see [11]. But the non-uniqueness will be still preserved. The category of this kind of representations is not semi-simple: there are non-trivial Ext's.

In [9, 10] the authors have developed harmonic analysis on the two-dimensional local fields and have defined the different types of functional spaces on these fields.

In [12, §5.4 (v)] the second named author asked a question on the structure of the natural representation of the discrete Heisenberg group $G = \mathrm{Heis}(3, \mathbb{Z})$ on the infinite-dimensional $\mathbb{C}$-vector space $V = \mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}$. Here K is a two-dimensional local field, $\mathcal{O}$ is the discrete valuation ring of the field K, $\mathcal{O}'$ is the ring of valuation of rank 2 of the field K, $\mathcal{D}'_{\mathcal{O}}(K)$ is the space of distributions on K introduced in [9] and [10].

In this paper we will give a partial answer to the above question. In particular, we will describe explicitly some irreducible subrepresentations of the representation of G on V and will relate these representations with classification of certain irreducible representation of G given in [11] and [1]. We note that the group $\tilde{G} = G \rtimes \mathbb{Z}$ also acts naturally on the space V. We will calculate the traces of the group $\tilde{G}$ on some irreducible subrepresentations of the representation of G on V. These traces are theta-series, as it was predicted in [11, Example] by another reasons.

Certainly, the results of this paper are only the first steps in the representation theory of discrete nilpotent groups. Recently, it was shown that a reasonable class of irreducible representations of any finitely generated nilpotent group coincides with the class of monomial representations, see [2]. This was known earlier for finite groups, Abelian groups and nilpotent groups of class 2. The general result was conjectured by the second named author in [12].

Acknowledgements. The first named author is grateful to the Institut des Hautes Études Scientifiques for the excellent working conditions, since the part of this research was done during his visit to IHÉS in October-November 2013.

2 Case of one-dimensional local fields

Let L be a one-dimensional local field, i.e. a complete discrete valuation field with a finite residue field $\mathbb{F}_q$. It is well-known that $L \simeq \mathbb{F}_q((u))$, or L is a finite extension of $\mathbb{Q}_p$. Let $\mathcal{O}_L$ be the discrete valuation ring of L, and $\mathfrak{m}_L$ be the maximal ideal of $\mathcal{O}_L$.

Let $\mathcal{D}(L)$ be the space of $\mathbb{C}$-valued locally constant functions with compact support on L, $\mathcal{E}(L)$ the space of $\mathbb{C}$-valued locally constant functions on L, and $\mathcal{D}'(L)$ be the space of distributions on L, i.e. the dual vector space to the discrete space $\mathcal{D}(L)$.

The additive group of the field L acts on the above spaces by translations. Let $\mu(L) = \mathcal{D}'(L)^L$ be the space of invariant elements under this action. In other words, it is the space of $\mathbb{C}$-valued Haar measures on L. We have $\dim_{\mathbb{C}}(\mu(L)) = 1$.

The group L^* acts on the additive group of the field L by multiplications. Hence we have an action of the group L^* on the the spaces $\mathcal{D}(L)$, $\mathcal{E}(L)$, $\mathcal{D}'(L)$, $\mu(L)$. We note that the subgroup $\mathcal{O}_L^*$ acts trivially on the space $\mu(L)$. Hence the group $\mathbb{Z} \simeq L^*/\mathcal{O}_L^*$ acts on the space $\mu(L)$ by the character $\mathbb{Z} \ni a \longmapsto q^a \in \mathbb{C}^*$. (A local parameter u from L^* goes to the element 1 from $\mathbb{Z}$ via the isomorphism $L^*/\mathcal{O}_L^* \simeq \mathbb{Z}$.)

We note that the group $\mathbb{Z} \simeq L^*/\mathcal{O}_L^*$ acts naturally on $\mathbb{C}$-vector spaces of invariant elements: $\mathcal{D}(L)^{\mathcal{O}_L^*}$, $\mathcal{E}(L)^{\mathcal{O}_L^*}$, $\mathcal{D}'(L)^{\mathcal{O}_L^*}$. We have also the following subrepresentations of the representation of the group $\mathbb{Z}$ on the space $\mathcal{D}'(L)^{\mathcal{O}_L^*}$:

$$\mathcal{D}(L)^{\mathcal{O}_L^*} \otimes_{\mathbb{C}} \mu(L) \ \subset \ \mathcal{E}(L)^{\mathcal{O}_L^*} \otimes_{\mathbb{C}} \mu(L) \ \subset \ \mathcal{D}'(L)^{\mathcal{O}_L^*}, \tag{1}$$

where the last embedding comes from the map:

$$f \otimes \nu \longmapsto \{g \mapsto \nu(fg)\} \qquad \forall \quad f \in \mathcal{E}(L), \quad g \in \mathcal{D}(L), \quad \nu \in \mu(L).$$

An explicit description of the space $\mathcal{D}(L)^{\mathcal{O}_L^*}$ as the space of some sequences is the following:

$$\mathcal{D}(L)^{\mathcal{O}_L^*} = \left\{ (c_n)_{n\in\mathbb{Z}} \;\middle|\; \begin{array}{l} c_n \in \mathbb{C}, \quad c_n = 0 \quad \text{if} \quad n \ll 0, \\ c_n = c_{n+1} = c_{n+2} = \ldots \quad \text{if} \quad n \gg 0 \end{array} \right\},$$

where by $f \in \mathcal{D}(L)^{\mathcal{O}_L^*}$ we construct $c_n = f(u^n) \in \mathbb{C}$ for $n \in \mathbb{Z}$ (here $u \in \mathcal{O}_L$ is a local parameter of the field L).

Proposition 1. *The representation of the group $\mathbb{Z}$ on the space $\mathcal{D}(L)^{\mathcal{O}_L^*}$ does not contain irreducible subrepresentations.*

Proof. We prove that for any nonzero $v \in \mathcal{D}(L)^{\mathcal{O}^*}$ the space $\mathbb{C}[\mathbb{Z}] \cdot v$ contains a proper nonzero $\mathbb{Z}$-invariant subspace. We consider an isomorphism:

$$\mathcal{D}(L)^{\mathcal{O}_L^*} \longrightarrow \mathbb{C}[z, z^{-1}] \qquad \text{given as} \qquad (c_n)_{n\in\mathbb{Z}} \longmapsto (1-z)(\sum_{n\in\mathbb{Z}} c_n z^n).$$

Under this isomorphism the action of an element $1 \in \mathbb{Z}$ goes to the multiplication on z. Therefore for any element $w \in \mathbb{C}[z, z^{-1}]$ we have $\mathbb{C}[\mathbb{Z}] \cdot w = \mathbb{C}[z, z^{-1}]w$. Now for any element $f \in \mathbb{C}[z]$ such that $f \notin \mathbb{C} \cdot z^k$ (for any integer k) we have that $\mathbb{C}[\mathbb{Z}] \cdot fw = \mathbb{C}[z, z^{-1}]fw$ is a proper nonzero $\mathbb{Z}$-invariant subspace in the space $\mathbb{C}[z, z^{-1}]w$. The proposition is proved. □

We note that between the spaces $\mathcal{D}(L)$ and $\mathcal{E}(L)$ there is a space $\tilde{\mathcal{E}}(L)$ that is the space of uniformly locally constant functions on L, i.e. f from $\mathcal{E}(L)$ belongs to $\tilde{\mathcal{E}}(L)$ if and only if there is $n_f \in \mathbb{N}$ such that $f(x+y) = f(x)$ for any $x \in L$, $y \in \mathfrak{m}_L^{n_f}$. It is clear that $\tilde{\mathcal{E}}(L)^{\mathcal{O}_L^*} = \mathcal{E}(L)^{\mathcal{O}_L^*}$ and

$$\mathcal{E}(L)^{\mathcal{O}_L^*} = \left\{ (c_n)_{n\in\mathbb{Z}} \;\middle|\; c_n \in \mathbb{C}, \quad , \quad c_n = c_{n+1} = c_{n+2} = \dots \quad \text{if} \quad n \gg 0 \right\}.$$

The one-dimensional subspace of all constant functions on L is the set of all eigenvectors in $\mathcal{E}(L)^{\mathcal{O}_L^*}$ with respect to the action of $1 \in \mathbb{Z}$.

We note that

$$\mathcal{D}(L) = \varinjlim_{m\in\mathbb{Z}} \mathcal{F}_0(m),$$

where $\mathcal{F}_0(m)$ is the space of all $\mathbb{C}$-valued functions with finite support on $L/\mathfrak{m}_L^m$. The transition map $\mathcal{F}_0(m_1) \to \mathcal{F}_0(m_2)$ for $m_1 < m_2$ is induced by the natural map $L/\mathfrak{m}_L^{m_2} \to L/\mathfrak{m}_L^{m_1}$, and the map $\mathcal{F}_0(m) \to \mathcal{D}(L)$ is induced by the map $L \to L/\mathfrak{m}_L^m$. Hence we have an equality

$$\mathcal{D}'(L) = \varprojlim_{m\in\mathbb{Z}} \mathcal{F}(m),$$

where $\mathcal{F}(m)$ is the space of all $\mathbb{C}$-valued functions on $L/\mathfrak{m}_L^m$. The transition map $\mathcal{F}(m_2) \to \mathcal{F}(m_1)$ for $m_1 < m_2$ is given by the sums of a function over the elements of finite fibers of the map $L/\mathfrak{m}_L^{m_2} \to L/\mathfrak{m}_L^{m_1}$, and the pairing $\mathcal{D}(L) \times \mathcal{D}'(L) \longrightarrow \mathbb{C}$ is induced by the pairing $\mathcal{F}_0(m) \times \mathcal{F}(m) \to \mathbb{C}$ for any integer m. The last pairing is given by the sum of product of two functions over elements of $L/\mathfrak{m}_L^m$. In a slightly another language this presentation of $D'(L)$ is described, for example, in [9, § 4.2]).

Now we can explicitly describe the space $\mathcal{D}'(L)^{\mathcal{O}_L^*}$:

$$\mathcal{D}'(L)^{\mathcal{O}_L^*} = \varprojlim_{m\in\mathbb{Z}} S(m), \tag{2}$$

where $S(m)$ is the space of sequences:

$$S(m) = \{ (a_n)_{n\le m} \mid a_n \in \mathbb{C} \}.$$

Any $f \in \mathcal{F}(m)^{\mathcal{O}_L^*}$ goes to $(a_n)_{n\le m} \in S(m)$, where $a_n = f(u^m)$ and $u \in \mathfrak{m}_L$ is a local parameter. The surjective maps $S(m+1) \to S(m)$ in the projective limit of formula (2) are the following:

$$(a_n)_{n\le m+1} \longmapsto (b_n)_{n\le m},$$

where $b_m = (q-1)a_m + a_{m+1}$, $b_{m-k} = qa_{m-k}$ if $k > 0$.

The pairing $\mathcal{D}(L) \times \mathcal{D}'(L) \longrightarrow \mathbb{C}$ induces the non-degenerate pairing

$$\mathcal{D}(L)^{\mathcal{O}_L^*} \times \mathcal{D}'(L)^{\mathcal{O}_L^*} \longrightarrow \mathbb{C}, \tag{3}$$

which can be described explicitly in the following way. We fix an element $(c_n)_{n\in\mathbb{Z}} \in \mathcal{D}(L)^{\mathcal{O}_L^*}$ and elements $(a_n)_{n\le m} \in S(m)$, $m \in \mathbb{Z}$ compatible with projective system (2). We choose any integers $N < M$ such that $c_n = 0$ for any $n < N$ and

$c_{M+k} = c_M$ for any $k > 0$. We define $l = M - N$. Now pairing (3) applied to elements $(c_n)_{n\in\mathbb{Z}}$ and $(a_n)_{n\le M} \in S(M)$ is equal to

$$(q^l - q^{l-1})c_N a_N + (q^{l-1} - q^{l-2})c_{N+1}a_{N+1} + \ldots + (q-1)c_{M-1}a_{M-1} + c_M a_M.$$

This result does not depend on the choice of appropriate M and N.

We note that the action of $1 \in \mathbb{Z}$ on $\mathcal{D}'(L)$ maps an element $(a_n)_{n\le m} \in S(m)$, $m \in \mathbb{Z}$ to an element $(b_n)_{n\le m+1} \in S(m+1)$, $m \in \mathbb{Z}$ where $b_n = a_{n-1}$ for $n \le m+1$.

Using explicit description of the space $\mathcal{D}'(L)^{\mathcal{O}_L^*}$ by formula (2), we obtain a proposition by direct calculations.

Proposition 2. *We fix any $\alpha \in \mathbb{C}^*$. We define a $\mathbb{C}$-vector space*

$$\Phi_\alpha = \left\{ \phi \in \mathcal{D}'(L)^{\mathcal{O}_L^*} \mid u(\phi) = \alpha\phi \right\},$$

where $u \in L^*$ is a local parameter, $\phi \longmapsto u(\phi) \in \mathcal{D}'(L)^{\mathcal{O}_L^*}$ is an action of $u \in L^*$ on ϕ (which is the same as an action of $1 \in \mathbb{Z}$ on ϕ). We have

$$\dim_{\mathbb{C}} \Phi_\alpha = 1.$$

Remark 1. For any $\alpha \in \mathbb{C}^*$ we can construct an element g_α from Φ_α explicitly:

$$(a_n)_{n\le m} \in S(m), \quad \text{where} \quad a_m = \alpha^{-m}, \quad a_{m-k} = \frac{\alpha^{-m+k-1}(\alpha-1)}{q^{k-1}(q-1)} \quad \text{for} \quad k > 0.$$

We note that the element g_α is uniquely defined inside the 1-dimensional space Φ_α by a property that the pairing of g_α with the characteristic function of the subgroup $\mathcal{O}_L$ is equal to 1.

Example 1. If $\alpha = 1$ then $\Phi_\alpha = \mathbb{C} \cdot \delta_0$, where δ_0 is the delta-function, i.e. $\delta_0(f) = f(0)$ for any $f \in \mathcal{D}(L)$. Besides, $g_1 = \delta_0$.

If $\alpha = q$, then $\Phi_\alpha = \mu(L)$ is the space of $\mathbb{C}$-valued Haar measures on L. Besides, g_q is the Haar measure which is equal to 1 after its application to the open compact subgroup $\mathcal{O}_L$.

Remark 2. The issues of this section were for the first time formulated and developed by A. Weil in [14] (see also a discussion in [13, Weil problem]).

3 Case of two-dimensional local fields

3.1 Two-dimensional local fields and the discrete Heisenberg group

Let K be a two-dimensional local field. Assume that K is isomorphic to one of the following fields (we will not consider another types of two-dimensional local fields):

$$\mathbb{F}_q((u))((t)), \qquad L((t)), \qquad M\{\{u\}\}, \tag{4}$$

where the fields L and M are finite extensions of the field $\mathbb{Q}_p$ such that the finite field $\mathbb{F}_q$ is the residue field of both L and M. (We recall that the field $M\{\{u\}\}$ is the completion of the field $\mathrm{Frac}(\mathcal{O}_M[[u]])$ with respect to the discrete valuation given by the prime ideal $\mathfrak{m}_M[[u]]$ of height 1 in the ring $\mathcal{O}_M[[u]]$. Here $\mathfrak{m}_M$ is the maximal ideal of the discrete valuation ring $\mathcal{O}_M$ of the field M.) See also, for example, [7, § 2] for the brief description how above types of two-dimensional local fields are constructed from an algebraic surface over a finite field or from an arithmetic surface.

Let u be a local parameter of L (in the second case of (4)), t be a local parameter of M (in the third case of (4)).

The field K is a discrete valuation field such that t is a local parameter of this valuation. Let $\mathcal{O}$ be the discrete valuation ring of K. We define the $\mathcal{O}$-module $\mathcal{O}(i) \subset K$ for any $i \in \mathbb{Z}$ as $\mathcal{O}(i) = t^i O$. We have $\mathcal{O}(0) = \mathcal{O}$, $\mathcal{O}(0)/\mathcal{O}(1)$ is the residue field of K. The image of the element u in the residue field is the local parameter there.

The pair t, u is called a system of local parameters of the field K.

We define a subring $\mathcal{O}' \subset \mathcal{O}$. We say that an element g from $\mathcal{O}$ belongs to $\mathcal{O}'$ if and only if the image of g in the residue field of K belongs to the discrete valuation ring of the residue field of K.

The field K has a valuation ν of rank 2 such that $\nu(f) = (a, b) \in \mathbb{Z}^2$ for $f \in K^*$ if and only if $f = t^a u^b g$ where $g \in \mathcal{O}'^*$. It is clear that the ring $\mathcal{O}'$ is the valuation ring of ν.

In accordance with (4) we define a subring B of K as one of the following expressions:

$$\mathbb{F}_q[[u]]((t)), \qquad \mathcal{O}_L((t)), \qquad M \cdot \mathcal{O}_M[[u]],$$

where $\mathcal{O}_L$ is the discrete valuation ring of L. (We note that the subring B depends on the choice of isomorphisms (4).)

Let $i \le j \in \mathbb{Z}$. An $\mathcal{O}$-module $\mathcal{O}(i)/\mathcal{O}(j)$ is a locally compact group with the base of neighbourhoods of 0 given by subgroups $(u^k B \cap \mathcal{O}(i))/(u^k B \cap \mathcal{O}(j))$, $k \in \mathbb{Z}$. For any $k \in \mathbb{Z}$ the subgroup $(u^k B \cap \mathcal{O}(i))/(u^k B \cap \mathcal{O}(j))$ is an open compact subgroup in $\mathcal{O}(i)/\mathcal{O}(j)$. We define a 1-dimensional $\mathbb{C}$-vector space $\mu(\mathcal{O}(i)/\mathcal{O}(j))$ as the space of $\mathbb{C}$-valued Haar measures on the group $\mathcal{O}(i)/\mathcal{O}(j)$. For any $r, s \in \mathbb{Z}$ we define a 1-dimensional $\mathbb{C}$-vector space $\mu(\mathcal{O}(r) \mid \mathcal{O}(s)) = \mu(\mathcal{O}(s)/\mathcal{O}(r))$ when $r \ge s$, and $\mu(\mathcal{O}(r) \mid \mathcal{O}(s)) = \mu(\mathcal{O}(r)/\mathcal{O}(s))^*$ when $r \le s$. We have that $\mu(\mathcal{O}(r) \mid \mathcal{O}(r))$ is canonically isomorphic to $\mathbb{C}$. There is also a canonical isomorphism for any elements r, s, w from $\mathbb{Z}$:

$$\mu(\mathcal{O}(r) \mid \mathcal{O}(s)) \otimes_{\mathbb{C}} \mu(\mathcal{O}(s) \mid \mathcal{O}(w)) \longrightarrow \mu(\mathcal{O}(r) \mid \mathcal{O}(w)) \tag{5}$$

satisfying associativity for any four elements from $\mathbb{Z}$. Any element $g \in K^*$ maps the $\mathbb{C}$-vector space $\mu(\mathcal{O}(r) \mid \mathcal{O}(s))$ to the $\mathbb{C}$-vector space $\mu(g\mathcal{O}(r) \mid g\mathcal{O}(s))$.

Lemma 1. *The group $\mathcal{O}'^*$ acts by the trivial character on the 1-dimensional $\mathbb{C}$-vector space $\mu(\mathcal{O}(s)/\mathcal{O}(r))$ for any integers $s \le r$.*

Proof. The proof follows by induction on $r - s$ with the help of formula (5). The case $r - s = 1$ is trivial as the case of one-dimensional local fields. The lemma is proved. □

We describe now a central extension of K^* by $\mathbb{C}^*$:

$$1 \longrightarrow \mathbb{C}^* \longrightarrow \widehat{K^*} \longrightarrow K^* \longrightarrow 1, \tag{6}$$

where the group $\widehat{K^*}$ consists from the pairs (g, μ): $g \in K^*$, $\mu \in \mu(\mathcal{O} \mid g\mathcal{O}))$ and $\mu \neq 0$. The group law in $\widehat{K^*}$ is the following: $(g_1, \mu_1) \cdot (g_2, \mu_2) = (g_1 g_2, \mu_1 \otimes g_1(\mu_2))$.

Now for any $i \leq j \in \mathbb{Z}$ instead the $\mathbb{C}^*$-torsor $\mu(\mathcal{O}(i)/\mathcal{O}(j)) \setminus 0$ we consider a $\mathbb{Z}$-torsor which is a set of Haar measures on the group $\mathcal{O}(i)/\mathcal{O}(j)$ with the property that the value of a Haar measure μ on the open compact subgroup $(B \cap \mathcal{O}(i))/(B \cap \mathcal{O}(j))$ is equal to q^b for some $b \in \mathbb{Z}$ (an integer b depends on μ). Using these $\mathbb{Z}$-torsors, by the same construction as for central extension (6) we obtain a central extension of K^* by $\mathbb{Z}$:

$$0 \longrightarrow \mathbb{Z} \longrightarrow \widetilde{K^*} \longrightarrow K^* \longrightarrow 1. \tag{7}$$

Besides, we have an embedding of central extension (7) to central extension (6): an embedding on kernels $\mathbb{Z} \ni b \longmapsto q^b \in \mathbb{C}^*$ gives an embedding of groups $\widetilde{K^*} \hookrightarrow \widehat{K^*}$.

We note that central extensions (7) and (6) split canonically over the subgroup $\mathcal{O}^*$ of the group K^*: $g \mapsto (g, 1)$, where $1 \in \mathbb{C} = \mu(\mathcal{O} \mid g\mathcal{O})$ since $g\mathcal{O} = \mathcal{O}$. Therefore these central extensions split canonically over the subgroup $\mathcal{O}'^*$, i. e. we can consider the group $\mathcal{O}'^*$ as a subgroup of the group $\widetilde{K^*}$. It is easy to see from Lemma 1 that the subgroup $\mathcal{O}'^*$ is contained in the center of the group $\widehat{K^*}$, and therefore this subgroup is also contained in the center of the group $\widetilde{K^*}$.

For any integers r and s we define an element $\mu_{B,r,s} \in \mu(\mathcal{O}(r) \mid \mathcal{O}(s))$ as following. If $s \leq r$, then $\mu(\mathcal{O}(r) \mid \mathcal{O}(s)) = \mu(\mathcal{O}(s)/\mathcal{O}(r))$, and the element $\mu_{B,r,s}$ is a Haar measure on the group $\mathcal{O}(s)/\mathcal{O}(r)$ such that the value of $\mu_{B,r,s}$ on the open compact subgroup $(B \cap \mathcal{O}(s))/(B \cap \mathcal{O}(r))$ is equal to 1. If $s \geq r$, then $\mu(\mathcal{O}(r) \mid \mathcal{O}(s)) = \mu(\mathcal{O}(r)/\mathcal{O}(s))^*$, and the element $\mu_{B,r,s}$ is the dual element to the element $\mu_{B,s,r} \in \mu(\mathcal{O}(s) \mid \mathcal{O}(r)) = \mu(\mathcal{O}(r)/\mathcal{O}(s))$. We note that $\mu_{B,r,r} = 1 \in \mathbb{C}$, and for any integers r, s and w the element $\mu_{B,r,s} \otimes \mu_{B,s,w}$ goes to the element $\mu_{B,r,w}$ via isomorphism (5).

We consider the discrete Heisenberg group $G = \mathrm{Heis}(3, \mathbb{Z})$ which is the group of matrices of the form:

$$\begin{pmatrix} 1 & a & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix},$$

where a, b, c are from $\mathbb{Z}$. We can consider also the group G as a set of all integer triples endowed with the group law:

$$(a_1, b_1, c_1)(a_2, b_2, c_2) = (a_1 + a_2, b_1 + b_2, c_1 + c_2 + a_1 b_2).$$

We note that G may be presented as

$$< \eta, \gamma \ : \ [\eta, [\eta, \gamma]] = 1 = [\gamma, [\eta, \gamma]] >$$

with η and γ corresponding to the generators $(1, 0, 0)$ and $(0, 1, 0)$ (see, for example, [5, § 2]). It is clear that the group G is nilpotent of class 2.

Proposition 3. *The group* $\widetilde{K^*}/\mathcal{O}'^*$ *is isomorphic to the group* $G = \mathrm{Heis}(3,\mathbb{Z})$.

Proof. We will construct this isomorphism explicitly. We consider a map

$$G \ni (a,b,c) \longmapsto (u^a t^b, q^{-c}\mu_{B,0,b}) \in \widetilde{K^*}. \tag{8}$$

Using the equality $B = tB$ we obtain an action of the element t as $t(\mu_{B,r,s}) = \mu_{B,r+1,s+1}$. Besides an action of the element u is $u(\mu_{B,r,s}) = q^{r-s}\mu_{B,r,s}$ (we recall that the pair t,u is the system of local parameters of K). Now from an explicit description of the group $\widetilde{K^*}$ given above, by a direct calculation we obtain that the composition of map (8) with the quotient map $\widetilde{K^*} \to \widetilde{K^*}/\mathcal{O}'^*$ is an isomorphism between groups G and $\widetilde{K^*}/\mathcal{O}'^*$. The proposition is proved. □

3.2 Representations of the discrete Heisenberg group

Let K be a two-dimensional local field that is isomorphic to one of the fields in formula (4). We keep the notation from Section 3.1.

The field K is a discrete valuation field. Its residue field $\bar{K} = \mathcal{O}/\mathcal{O}(1)$ is isomorphic to the following field (according to formula (4)):

$$\mathbb{F}_q((u)), \qquad L, \qquad \mathbb{F}_q((u)).$$

For any integers $s \le r$ the group $\mathcal{O}(s)/\mathcal{O}(r)$ is a locally compact totally disconnected group. Let $\mathcal{D}(\mathcal{O}(s)/\mathcal{O}(r))$ be the space of $\mathbb{C}$-valued locally constant functions with compact support on the group $\mathcal{O}(s)/\mathcal{O}(r)$. Let $\mathcal{D}'(\mathcal{O}(s)/\mathcal{O}(r))$ be the space of distributions on the group $\mathcal{O}(s)/\mathcal{O}(r)$ (i.e. the dual vector space to the discrete space $\mathcal{D}(\mathcal{O}(s)/\mathcal{O}(r))$).

The following spaces which are two-dimensional analogs of the classical function and distribution spaces were constructed in [9, § 5.3] and [10, § 7] using an idea from [6]:

$$\mathcal{D}_{\mathcal{O}}(K) = \varprojlim_{s} \varprojlim_{r>s} \mathcal{D}(\mathcal{O}(s)/\mathcal{O}(r)) \otimes_{\mathbb{C}} \mu(\mathcal{O}(r) \mid \mathcal{O}), \tag{9}$$

$$\mathcal{D}'_{\mathcal{O}}(K) = \varinjlim_{s} \varinjlim_{r>s} \mathcal{D}'(\mathcal{O}(s)/\mathcal{O}(r)) \otimes_{\mathbb{C}} \mu(\mathcal{O} \mid \mathcal{O}(r)), \tag{10}$$

where transition maps in above limits are given by direct and inverse images of function and distribution spaces on locally compact groups.

There is a natural non-degenerate pairing:

$$\mathcal{D}_{\mathcal{O}}(K) \times \mathcal{D}'_{\mathcal{O}}(K) \longrightarrow \mathbb{C}.$$

The group $\widehat{K^*}$ acts naturally on the spaces $\mathcal{D}_{\mathcal{O}}(K)$ and $\mathcal{D}'_{\mathcal{O}}(K)$. We recall this action on $\mathcal{D}'_{\mathcal{O}}(K)$ (the case of $\mathcal{D}_{\mathcal{O}}(K)$ is analogous). We have to apply limits to the following action of an element (g,μ) from $\widehat{K^*}$:

$$\mathcal{D}'(\mathcal{O}(s)/\mathcal{O}(r)) \otimes_{\mathbb{C}} \mu(\mathcal{O} \mid \mathcal{O}(r)) \xrightarrow{g(\,)} \mathcal{D}'(g\mathcal{O}(s)/g\mathcal{O}(r)) \otimes_{\mathbb{C}} \mu(g\mathcal{O} \mid g\mathcal{O}(r))$$

$$\mathcal{D}'(g\mathcal{O}(s)/g\mathcal{O}(r)) \otimes_{\mathbb{C}} \mu(g\mathcal{O} \mid g\mathcal{O}(r)) \xrightarrow{\mu\otimes} \mathcal{D}'(g\mathcal{O}(s)/g\mathcal{O}(r)) \otimes_{\mathbb{C}} \mu(\mathcal{O} \mid g\mathcal{O}(r)).$$

We recall that the group $\mathcal{O}'^*$ can be considered as a subgroup of the group $\widehat{K^*}$. We are interested in the spaces of invariant elements with respect to the action of the group $\mathcal{O}'^*$: $\mathcal{D}_{\mathcal{O}}(K)^{\mathcal{O}'^*}$ and $\mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}$.

Proposition 4. *We have* $\mathcal{D}_{\mathcal{O}}(K)^{\mathcal{O}'^*} = 0$.

Proof. We have Lemma 1. Besides, the transition maps in projective limits in formula (9) are $\mathcal{O}'^*$-equivariant maps. Therefore it is enough to prove that $\mathcal{D}(\mathcal{O}(s)/\mathcal{O}(r))^{\mathcal{O}'^*} = 0$ for $r - s > 1$. Assume the converse. We take a non-zero element $f \in \mathcal{D}(\mathcal{O}(s)/\mathcal{O}(r))^{\mathcal{O}'^*}$. Since f has a compact support, there is an integer l such that $f(x) = 0$ when $x \notin u^l V$, where $V = (B \cap \mathcal{O}(s))/(B \cap \mathcal{O}(r))$. It contradicts to the fact $f(gy) = f(y)$ where g is any from $\mathcal{O}'^*$, y is any from $\mathcal{O}(s)/\mathcal{O}(r)$, because if $f(y) \neq 0$, then there is $g \in \mathcal{O}'^*$ such that $gy \notin u^l V$, and hence $f(gy) \neq 0$. The proposition is proved. □

The group $G = \widetilde{K^*}/\mathcal{O}'^*$ acts naturally on the space $\mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}$.

Problem. How to describe the space $\mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}$ and the action of the group G on it? (See also [12, §5.4 (v)].)

We will describe the following G-invariant infinite-dimensional subspace of the space $\mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}$.

Let $\varphi \in \mathcal{D}'(\bar{K})^{\mathcal{O}^*_{\bar{K}}} = \mathcal{D}'(\mathcal{O}/\mathcal{O}(1))^{\mathcal{O}^*_{\bar{K}}}$. For any integer l we *define* an element

$$\varphi_l = t^l(\varphi) \otimes \mu_{B,0,l+1} \ \in \ \mathcal{D}'_{\mathcal{O}}(K), \tag{11}$$

where an element $t^l(\varphi) \in \mathcal{D}'(\mathcal{O}(l)/\mathcal{O}(l+1))$ is an application of the action of t^l to φ, an element $\mu_{B,0,l+1} \in \mu(\mathcal{O} \mid \mathcal{O}(l+1))$ was defined in Section 3.1. We have that $\varphi_l \in \mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}$, because $\varphi \in \mathcal{D}'(\mathcal{O}/\mathcal{O}(1))^{\mathcal{O}^*_{\bar{K}}}$ and, by lemma 1, $g\mu_{B,0,l+1} = \mu_{B,0,l+1}$ for any $g \in \mathcal{O}'^*$, and the transition maps in inductive limits in formula (10) are $\mathcal{O}'^*$-equivariant maps.

Definition 1. *We define a* $\mathbb{C}$*-vector subspace* Ψ *of the vector space* $\mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}$ *as the* $\mathbb{C}$*-linear span of elements* φ_l *for all elements* $\varphi \in \mathcal{D}'(\bar{K})^{\mathcal{O}^*_{\bar{K}}}$ *and all integers* l.

Using en explicit isomorphism from the proof of Proposition 3 and a description of the action of $\widehat{K^*}$ on $\mathcal{D}'_{\mathcal{O}}(K)$, we obtain the following proposition.

Proposition 5. *The space* Ψ *is a* G*-invariant subspace with an explicit action:*

$$(a,b,c) \cdot \varphi_l = q^{-c-a(l+1)} a(\varphi)_{l+b},$$

where $(a,b,c) \in G$ *and* $a(\varphi) \in \mathcal{D}'(\bar{K})^{\mathcal{O}^*_{\bar{K}}}$ *is an application of the action of the element* $u^a \in \bar{K}^*$ *to the element* φ.

Proof. By direct calculations we have

$$\begin{aligned}(a,b,c) \cdot \varphi_l = \left(u^a t^b, q^{-c}\mu_{B,0,b}\right) \cdot \left(t^l(\varphi) \otimes \mu_{B,0,l+1}\right) = \\ = \left(t^{l+b}(u^a(\varphi)), q^{-c}\mu_{B,0,b} \otimes u^a(\mu_{B,b,b+l+1}\right) = \\ = \left(t^{l+b}(a(\varphi)), q^{-c-a(l+1)}\mu_{B,0,b} \otimes \mu_{B,b,b+l+1}\right) = q^{-c-a(l+1)} a(\varphi)_{l+b}.\end{aligned}$$

The proposition is proved.

□

We recall the notion of the induced representation for discrete groups.

Definition 2. *Let $G_1 \subset G_2$ be groups, and $\tau : G_1 \to \mathrm{Aut}_{\mathbb{C}} V$ be a representation of the group G_1 in a $\mathbb{C}$-vector space V. A space of the induced representation $\mathrm{ind}_{G_1}^{G_2}(\tau)$ consists of all functions $f : G_2 \to V$ subject to the following conditions:*

1) $f(g_2 g_1) = \tau(g_1^{-1}) f(g_2)$ for any elements $g_1 \in G_1$, $g_2 \in G_2$,

2) the support $\mathrm{Supp}(f)$ is contained in finite number of left cosets G_2/G_1.

An action of the group G_2 is the following: for any $f \in \mathrm{ind}_{G_1}^{G_2}(\tau)$ and $g \in G_2$ we have $(g \cdot f)(x) = f(g^{-1}x)$ where $x \in G_2$.

We consider subgroups $H = \mathbb{Z}$, $C = \mathbb{Z}$ and $P = \mathbb{Z}$ of the group G given as

$$H \ni a \mapsto (a, 0, 0) \in G, \qquad P \ni b \mapsto (0, b, 0) \in G, \qquad C \ni c \mapsto (0, 0, c),$$

We consider a representation

$$\sigma : H \oplus C \longrightarrow \mathrm{Aut}_{\mathbb{C}}(\mathcal{D}'(\bar{K})^{\mathcal{O}_{\bar{K}}^*}), \qquad \sigma(a \oplus c)(\varphi) = q^{-c-a} a(\varphi), \tag{12}$$

where $\varphi \in \mathcal{D}'(\bar{K})^{\mathcal{O}_{\bar{K}}^*}$ and $a(\varphi) = u^a(\varphi) \in \mathcal{D}'(\bar{K})^{\mathcal{O}_{\bar{K}}^*}$.

The group $H \oplus C$ is a subgroup of the group G via the map $a \oplus c \mapsto (a, 0, c)$. We note that the set of left cosets $G/(H \oplus C)$ can be naturally identified with the set of elements of the subgroup P. By Definition 2, the space $\mathrm{ind}_{H\oplus C}^{G}(\sigma)$ can be identified with the space of functions with finite support from P to $\mathcal{D}'(\bar{K})^{\mathcal{O}_{\bar{K}}^*}$. Therefore the space $\mathrm{ind}_{H\oplus C}^{G}(\sigma)$ is linearly generated by elements $\Delta(m, \varphi)$ where $m \in \mathbb{Z}$, $\varphi \in \mathcal{D}'(\bar{K})^{\mathcal{O}_{\bar{K}}^*}$. Here $\Delta(m, \varphi)$ is a function from $P = \mathbb{Z}$ to $\mathcal{D}'(\bar{K})^{\mathcal{O}_{\bar{K}}^*}$ such that

- $\Delta(m, \varphi)(n) = \varphi$ if $m = n$,
- $\Delta(m, \varphi)(n) = 0$ if $m \neq 0$.

We consider *a map* β from the space $\mathrm{ind}_{H\oplus C}^{G}(\sigma)$ to the space Ψ given as

$$\beta(\Delta(m, \varphi)) = \varphi_m.$$

Theorem 1. *The map β induces an exact sequence of G-representations*

$$0 \longrightarrow \mathrm{Ker}\, \beta \longrightarrow \mathrm{ind}_{H\oplus C}^{G}(\sigma) \xrightarrow{\beta} \Psi \longrightarrow 0.$$

The subspace $\mathrm{Ker}\, \beta$ has a basis which consists of elements

$$v_m = \Delta(m, \delta_0) - \Delta(m + 1, \mu_{B,1,0}) \qquad (m \in \mathbb{Z})$$

where δ_0 is the delta-function at 0, and $\mu_{B,1,0} \in \mu(\bar{K}) \subset \mathcal{D}'(\bar{K})^{\mathcal{O}_{\bar{K}}^*}$. Moreover, the space $\mathrm{Ker}\, \beta$ is an irreducible G-representation which is isomorphic to G-representation $\mathrm{ind}_{H\oplus C}^{G}(\chi_0)$ with the character $\chi_0 : H \oplus C \longrightarrow \mathbb{C}^*$ given as $\chi_0(a \oplus c) = q^{-c}$.

Proof. We check that the map β is a map of G-representations. Let $g = (a, b, c) \in G$. We have $g^{-1} = (-a, -b, ab - c)$. Using the first condition from definition 2 we extend a function $\Delta(m, \varphi)$ to a function from G to $\mathcal{D}'(\bar{K})^{\mathcal{O}_K^*}$. We have for any $g = (a, b, c)$ and any integer k

$$\begin{aligned}(g \cdot \Delta(m,\varphi))((0,k,0)) = \Delta(m,\varphi))(g^{-1} \cdot (0,k,0)) = \\ = \Delta(m,\varphi)((-a,-b,ab-c)(0,k,0)) = \Delta(m,\varphi)((-a,-b+k,ab-ak-c)) = \\ = \Delta(m,\varphi)((0,-b+k,0)(-a,0,ab-ak-c)) = \\ = \sigma\big(a \oplus (c+ak-ab)\big)\big(\Delta(m,\varphi)((0,-b+k,0))\big) = \\ = q^{-c-a(k-b+1)}a\big(\Delta(m,\varphi)((0,-b+k,0))\big). \quad (13)\end{aligned}$$

Now if $k = b + m$ then the last formula is equal to $q^{-c-a(m+1)}a(\varphi)$. If $k \neq b + m$, then the last formula is equal to 0. Hence we have

$$(a, b, c) \cdot \Delta(m, \varphi) = q^{-c-a(m+1)}\Delta(b + m, a(\varphi)). \quad (14)$$

Therefore using the formula from Proposition 5 we obtain that the map β commutes with the action of G.

The surjectivity of the map β follows from the definition of this map. We calculate the kernel of β. We have from the construction that elements v_m ($m \in \mathbb{Z}$) belong to the space $\operatorname{Ker}\beta$. It is also clear that elements v_m ($m \in \mathbb{Z}$) are linearly independent over $\mathbb{C}$. Let an element $\sum\limits_{i=1}^{n} \Delta(m_i, \varphi_i)$ belongs to $\operatorname{Ker}\beta$, where $m_i < m_{i+1}$ for $1 \leq i \leq (n-1)$, and $\varphi_i \in \mathcal{D}'(\bar{K})^{\mathcal{O}_K^*}$ for $1 \leq i \leq n$. Hence we obtain

$$\sum_{i=1}^{n}(\varphi_i)_{m_i} = 0 \quad (15)$$

in the space Ψ. Moreover, equality (15) is satisfied in the space

$$\mathcal{D}'(\mathcal{O}(m_1)/\mathcal{O}(m_n + 1)) \otimes_{\mathbb{C}} \mu(\mathcal{O}(m_n + 1) \mid \mathcal{O}),$$

since the transition maps in limits in formula (10) are injective. We fix $\mu_{B,m_n+1,0} \in \mu(\mathcal{O}(m_n + 1) \mid \mathcal{O})$. After that we can consider an analog of equality (15) in $\mathcal{D}'(\mathcal{O}(m_1)/\mathcal{O}(m_n+1))$. In the last equality we consider the supports of summands, where every summand is an element of $\mathcal{D}'(\mathcal{O}(m_1)/\mathcal{O}(m_n + 1))$ (see, for example, the definition of the support of distribution in [9, Def. 4]). From this equality we obtain that $\operatorname{supp}(\varphi_1) = \{0\}$ as a subset in $\bar{K} = \mathcal{O}/\mathcal{O}(1)$. Therefore $1(\varphi_1) = \varphi_1$, and from Example 1 we obtain that $(\varphi_1)_{m_1} = e \cdot (\delta_0)_{m_1}$, where $e \in \mathbb{C}$. Now we use $(\delta_0)_{m_1} = (\mu_{B,1,0})_{m_1+1}$. From induction on the length $m_{n+1} - m_1$ we obtain that the space $\operatorname{Ker}\beta$ is linearly generated by elements v_m with $m \in \mathbb{Z}$.

From formula (14) we have $(a, b, c) \cdot v_m = q^{-c-a(m+1)}v_{m+b}$ for any $(a, b, c) \in G$ and $m \in \mathbb{Z}$. Using calculations analogous to (13), it is easy to see that this action corresponds to the action of G on $\operatorname{ind}_{H\oplus C}^{G}(\chi)$ (if we consider elements v_m as functions δ_m from the subgroup P to $\mathbb{C}$ supported on $m \in \mathbb{Z}$), where the character χ is given as $\chi(a \oplus c) = q^{-c-a}$. If we shift the basis: $\tilde{v}_m = v_{m-1}$, then the action is

$$(a, b, c) \cdot \tilde{v}_m = q^{-c-am}\tilde{v}_{m+b}. \quad (16)$$

The last action corresponds to the action of G on $\mathrm{ind}_{H\oplus C}^{G}(\chi_0)$, where the character χ_0 is given as $\chi_0(a\oplus c)=q^{-c}$. Thus, G-representations $\mathrm{ind}_{H\oplus C}^{G}(\chi)$ and $\mathrm{ind}_{H\oplus C}^{G}(\chi_0)$ are isomorphic.

We explain the irreducibility of G-representation $\mathrm{ind}_{H\oplus C}^{G}(\chi_0)$ (see another argument in [11, Prop. 1]). Consider any nonzero element

$$x=\sum_{i=1}^{n}\alpha_i\tilde{v}_{m_i}\ \in\ \mathrm{Ker}\,\beta,$$

where $m_i\in\mathbb{Z}$ and $\alpha_i\in\mathbb{C}^*$ for $1\le i\le n$, besides $m_k<m_{k+1}$ for $1\le k\le n-1$. We will prove that $\mathbb{C}[G]\cdot x=\mathrm{Ker}\,\beta$. Consider a column $Y=[y_1;\ldots;y_n]$ whose entries $y_j=(-j+1,0,0)\cdot x$ with $1\le j\le n$ are elements of the space $\mathrm{Ker}\,\beta$. From formula (16) we obtain that $Y=AZ$, where the column $Z=[z_1;\ldots;z_n]$ has entries $z_i=\alpha_i\tilde{v}_{m_i}$, where $1\le i\le n$, and the entries of the square matrix A are as follows: $A_{j,i}=q^{(j-1)m_i}$, where $1\le j\le n$ and $1\le i\le n$. The matrix A is the Vandermonde matrix with non-zero determinant equal to

$$\prod_{1\le j_1<j_2\le n}(q^{m_{j_2}}-q^{m_{j_1}}).$$

Therefore there is the inverse matrix A^{-1} with entries from the field $\mathbb{Q}$. Hence we obtain $Z=A^{-1}AZ=A^{-1}Y$. Therefore any element $z_i=\alpha_i\tilde{v}_{m_i}$, where $1\le i\le n$, is a linear combination (with coefficients from $\mathbb{Q}$) of elements $y_j=(-j+1,0,0)\cdot x$, where $1\le j\le n$. Hence any element $\tilde{v}_{m_i}$, where $1\le i\le n$, belongs to the space $\mathbb{C}[G]\cdot x$. By formula (16), the action by means of an element $(0,b,0)$, where $b\in\mathbb{Z}$, maps an element $\tilde{v}_k$ to the element $\tilde{v}_{k+b}$, where $k\in\mathbb{Z}$. Therefore, we obtain that any element $\tilde{v}_i$ with $i\in\mathbb{Z}$ belongs to the space $\mathbb{C}[G]\cdot x$. The theorem is proved. □

We consider subrepresentations $\sigma_1\subset\sigma_2$ of the representation σ (see formula (12) and formula (1)):

$$\sigma_1: H\oplus C\longrightarrow \mathrm{Aut}_{\mathbb{C}}(\mathcal{D}(\bar{K})^{\mathcal{O}_{\bar{K}}^*}\otimes_{\mathbb{C}}\mu(\bar{K})),$$

$$\sigma_2: H\oplus C\longrightarrow \mathrm{Aut}_{\mathbb{C}}(\mathcal{E}(\bar{K})^{\mathcal{O}_{\bar{K}}^*}\otimes_{\mathbb{C}}\mu(\bar{K})).$$

We have the following embeddings of representations of the group G:

$$\mathrm{ind}_{H\oplus C}^{G}(\sigma_1)\ \subset\ \mathrm{ind}_{H\oplus C}^{G}(\sigma_2)\ \subset\ \mathrm{ind}_{H\oplus C}^{G}(\sigma).$$

Proposition 6. *The map β restricted to $\mathrm{ind}_{H\oplus C}^{G}(\sigma_2)$ is an isomorphism with its image in Ψ.*

Proof. From theorem 1 we have an explicit description of elements v_m ($m\in\mathbb{Z}$) which form a basis in the space $\mathrm{Ker}\,\beta$. Since the element δ_0 from the space $\mathcal{D}'(\bar{K})$ does not belong to the subspace $\mathcal{E}(\bar{K})\otimes_{\mathbb{C}}\mu(\bar{K})$, we obtain that any non-zero element $\sum_{i=1}^{n}a_iv_i$ ($a_i\in\mathbb{C}$) does not belong to $\mathrm{ind}_{H\oplus C}^{G}(\sigma_2)$. The proposition is proved. □

Let $\alpha \in \mathbb{C}^*$. We recall that the 1-dimensional $\mathbb{C}$-vector subspace Φ_α of the space $\mathcal{D}'(\bar{K})^{\mathcal{O}_{\bar{K}}^*}$ was introduced in Proposition 2. We consider a subrepresentation σ_α of the representation σ (see formula (12)):

$$\sigma_\alpha : H \oplus C \longrightarrow \mathrm{Aut}_{\mathbb{C}}(\Phi_\alpha) = \mathbb{C}^*, \qquad \sigma_\alpha(a \oplus c) \longmapsto q^{-c-a}\alpha^a.$$

We define a G-subrepresentation $\Psi_\alpha = \mathrm{ind}_{H\oplus C}^G(\sigma_\alpha)$ of G-representation $\mathrm{ind}_{H\oplus C}^G(\sigma)$. We consider the element $g_\alpha \in \Phi_\alpha$ (see Remark 1). Now elements $\Delta(m, g_\alpha)$ ($m \in \mathbb{Z}$) form a basis of the subspace Ψ_α inside the space $\mathrm{ind}_{H\oplus C}^G(\sigma)$. From formula (14) we have the following action of the group G on these elements:

$$(a, b, c) \cdot \Delta(m, g_\alpha) = q^{-c-a(m+1)}\alpha^a \Delta(b + m, g_\alpha). \tag{17}$$

Theorem 2. *G-representations Ψ_α have the following properties.*

1. *For any $\alpha \in \mathbb{C}^*$ the space Ψ_α is an irreducible G-representation, and the map β restricted to Ψ_α is an isomorphism with its image in Ψ.*
2. *If $\alpha = q$, then $\Psi_\alpha \subset \mathrm{ind}_{H\oplus C}^G(\sigma_2)$. If $\alpha \neq q$, then $\Psi_\alpha \cap \mathrm{ind}_{H\oplus C}^G(\sigma_2) = \{0\}$ inside $\mathrm{ind}_{H\oplus C}^G(\sigma)$.*
3. *We have $\beta(\Psi_1) = \beta(\Psi_q)$ as subspaces in the space Ψ. For any pair $(\alpha_1, \alpha_2) \in \mathbb{C}^* \times \mathbb{C}^*$ such that $\alpha_1 \neq \alpha_2$, $(\alpha_1, \alpha_2) \neq (q, 1)$ and $(\alpha_1, \alpha_2) \neq (1, q)$ we have $\beta(\Psi_{\alpha_1}) \cap \beta(\Psi_{\alpha_2}) = \{0\}$ inside the space Ψ.*
4. *We have embeddings of G-representations:*

$$\bigoplus_{\alpha\in\mathbb{C}^*} \Psi_\alpha \subset \mathrm{ind}_{H\oplus C}^G(\sigma) \qquad \text{and} \qquad \bigoplus_{\substack{\alpha\in\mathbb{C}^* \\ \alpha\neq 1}} \beta(\Psi_\alpha) \subset \Psi. \tag{18}$$

5. *For any α_1 and α_2 from $\mathbb{C}^*$ we have that the G-representations Ψ_{α_1} and Ψ_{α_2} are isomorphic if and only if $\alpha_1 = q^l\alpha_2$ for some integer l. Thus the G-representation Ψ contains irreducible G-representations which are parametrized by the elliptic curve $\mathbb{C}^*/q^{\mathbb{Z}}$. Any such irreducible G-representation is included with at least countable multiplicity in the G-representation Ψ.*

Proof. 1. The space Ψ_α is an irreducible G-representation by an argument which is similar to the end of the proof of Theorem 1 (see another argument in [11, Prop. 1]). We prove that $\mathrm{Ker}\,\beta \cap \Psi_\alpha = 0$. By Theorem 1 the G-representation $\mathrm{Ker}\,\beta$ is irreducible. Therefore it is enough to prove that an element v_m from $\mathrm{Ker}\,\beta$ does not belong to Ψ_α. The latter follows from the fact that $\delta_0 \in \Phi_0$ and $\mu_{B,1,0} \in \Phi_q$ (see Example 1).

2. We have $\Psi_q \subset \mathrm{ind}_{H\oplus C}^G(\sigma_2)$, since the space Φ_q is equal to the space $\mu(\bar{K})$ of $\mathbb{C}$-valued Haar measures on $\bar{K}$ (see Example 1). The subspace of all constant functions on $\bar{K}$ is the set of all eigenvectors in $\mathcal{E}(\bar{K})^{\mathcal{O}_{\bar{K}^*}}$ with respect to the action of local parameter $u \in \bar{K}^*$ (or $1 \in \mathbb{Z}$). Therefore if $\alpha \neq q$, then $\Phi_\alpha \cap (\mathcal{E}(\bar{K})^{\mathcal{O}_{\bar{K}}^*} \otimes_{\mathbb{C}} \mu(\bar{K})) = \{0\}$. Hence if $\alpha \neq q$, then $\Psi_\alpha \cap \mathrm{ind}_{H\oplus C}^G(\sigma_2) = \{0\}$.

3. This item follows from explicit description of basis vectors v_m ($m \in \mathbb{Z}$) of the space $\mathrm{Ker}\,\beta$ (see Theorem 1), from description of basis vectors of the space Ψ_α , and from assertion 1.

4. Elements $\Delta(m, g_\alpha)$ ($m \in \mathbb{Z}$, $\alpha \in \mathbb{C}^*$) from $\mathrm{ind}_{H\oplus C}^{G}(\sigma)$ are linearly independent over $\mathbb{C}$. (Indeed, it is enough to see that elements g_α with different $\alpha \in \mathbb{C}^*$ are linearly independent in $\mathcal{D}'(\bar{K})$. This is true, because these elements are eigenvectors with different eigenvalues of linear operator induced by multiplication action of local parameter $u \in \bar{K}^*$.) These elements form a basis of the space $\bigoplus\limits_{\alpha\in\mathbb{C}^*} \Psi_\alpha$. Thus we obtained the first formula in (18). We note that

$$\mathrm{Ker}\,\beta \ \subset\ (\Psi_1 \oplus \Psi_q) \qquad \text{and} \qquad (\Psi_1 \oplus \Psi_q)/\mathrm{Ker}\,\beta \ =\ \beta(\Psi_q).$$

   Hence and from the first formula in (18) we obtain the second formula in (18).

5. For this item we use the previous items of the theorem. Now we prove that G-representations Ψ_{α_1} and Ψ_{α_2} are isomorphic if and only if $\alpha_1 = q^l\alpha_2$ for some integer l (see also [11, Prop. 1] and [1, Th. 5]). If $\alpha_1 = q^l\alpha_2$, then this isomorphism is constructed by a shift of the elements of the basis: $\Delta(m, g_{\alpha_1}) \mapsto \Delta(m - l, g_{\alpha_2})$ (see formula (17)). Suppose that above two representations are isomorphic. We note that elements of the one-dimensional vector spaces $\mathbb{C}\cdot\Delta(m, g_{\alpha_i})$ are the only eigenvectors of elements of the subgroup H under the action on Ψ_{α_i} . (In other words, we have the weight decomposition of Ψ_{α_i} in above one-dimensional vector spaces with respect to the subgroup H .) Therefore the isomorphism of G-representations maps the element $\Delta(0, g_{\alpha_1})$ to the element $d\Delta(-l, g_{\alpha_2})$ for certain integer l and $d \in \mathbb{C}^*$. The action of the element $(1, 0, 0)$ from G on above two elements by formula (17) gives $\alpha_1 = q^l\alpha_2$.

   The theorem is proved.

□

4 Fourier transform

Let L be a one-dimensional local field with a finite residue field $\mathbb{F}_q$ (see Section 2). Let u be a local parameter in L . Then the additive group of the field L is identified with its dual Pontryagin group via the pairing:

$$\chi\ :\ L \times L \longrightarrow \mathbb{T}, \qquad\qquad \chi(l_1, l_2) = \exp(2\pi i \cdot \lambda_L(l_1 l_2)),$$

where $\mathbb{T}$ is the circle group, i.e. the unit circle in $\mathbb{C}$, and the map $\lambda_L : L \to \mathbb{Q}/\mathbb{Z}$ is defined as following. If $L \simeq \mathbb{F}_q((u))$, then for an element $l = \sum a_i u^i$ from L (where $a_i \in \mathbb{F}_q$) we define $\lambda_L(l) = \mathrm{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(a_{-1}) \in p^{-1}\mathbb{Z}/\mathbb{Z}$. If L is a finite extension of $\mathbb{Q}_p$, then the map λ_L is the composition of maps:

$$L \ \stackrel{\mathrm{Tr}_{L/\mathbb{Q}_p}}{\longrightarrow}\ \mathbb{Q}_p \longrightarrow \mathbb{Q}_p/\mathbb{Z}_p \longrightarrow \mathbb{Q}/\mathbb{Z}.$$

For any $\mu \in \mu(L) \subset \mathcal{D}'(L)$, $\mu \neq 0$ we have the Fourier transform

$$\mathbf{F}_\mu\ :\ \mathcal{D}(L) \to \mathcal{D}(L), \qquad\qquad (\mathbf{F}_\mu(f))(y) = \mu(f \cdot \chi_{-y})),$$

where $f \in \mathcal{D}(L)$, $y \in L$, and χ_{-y} is a locally constant function on L defined as $\chi_{-y}(x) = \chi(-y, x)$ for $x \in L$. As the dual map to the above map we have the Fourier transform on the distribution space

$$\mathbf{F}_{\mu^{-1}} \ : \ \mathcal{D}'(L) \to \mathcal{D}'(L).$$

By direct calculations, for any $l \in L^*$ and $f \in \mathcal{D}(L)$ we have

$$\mathbf{F}_{\mu}(l(f)) = (\mu/l(\mu)) \cdot l^{-1}(\mathbf{F}_{\mu}(f)),$$

where $(\mu/l(\mu)) \in \mathbb{C}^*$. Hence for any $G \in \mathcal{D}'(L)$ we obtain

$$\mathbf{F}_{\mu^{-1}}(l(G)) = (l(\mu)/\mu) \cdot l^{-1}(\mathbf{F}_{\mu^{-1}}(G)).$$

Hence we have well-defined maps

$$\mathbf{F}_{\mu} : \mathcal{D}(L)^{\mathcal{O}_L^*} \longrightarrow \mathcal{D}(L)^{\mathcal{O}_L^*}, \qquad \mathbf{F}_{\mu^{-1}} : \mathcal{D}'(L)^{\mathcal{O}_L^*} \longrightarrow \mathcal{D}'(L)^{\mathcal{O}_L^*}$$

and the following proposition about spaces Φ_α (recall also Proposition 2).

Proposition 7. *For any $\alpha \in \mathbb{C}^*$ we have $\mathbf{F}_{\mu^{-1}}(\Phi_\alpha) = \Phi_{q\alpha^{-1}}$.*

Remark 3. We recall that in Remark 1 we have constructed elements $g_\alpha \in \Phi_\alpha$ for any $\alpha \in \mathbb{C}^*$. Let μ_0 is a Haar measure on L such that its appication to the open compact subgroup $\mathcal{O}_L$ is equal to 1. We have $\mathbf{F}_{\mu_0}(g_\alpha) = g_{q\alpha^{-1}}$.

Let K be a two-dimensional local field that is isomorphic to one of the fields in formula (4). We keep the notation from Sections 3.1 and 3.2. The additive group of the field K is self-dual via the pairing:

$$K \times K \longrightarrow \mathbb{T}, \qquad k_1 \times k_2 \longmapsto \exp(2\pi i \cdot \Lambda(k_1 \cdot k_2)),$$

where the map $\Lambda : K \longrightarrow \mathbb{Q}/\mathbb{Z}$ is defined as following. If K is the first field from formula (4), then for an element $k = \sum a_{i,j} u^i t^j$ from K (where $a_{i,j} \in \mathbb{F}_q$) we define $\Lambda(k) = \mathrm{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(a_{-1,-1}) \in p^{-1}\mathbb{Z}/\mathbb{Z}$. If K is the second field from formula (4), then for an element $k = \sum a_i t^i$ from K (where $a_i \in L$) we define $\Lambda(k) = \lambda_L(a_{-1}) \in \mathbb{Q}/\mathbb{Z}$. If K is the third field from formula (4), then an element k from K can be presented as infinite in both sides sum $\sum a_i u^i$ (where $a_i \in M$), and, by definition, $\Lambda(k) = \lambda_M(a_{-1}) \in \mathbb{Q}/\mathbb{Z}$.

The two-dimensional Fourier transform

$$\mathbf{F} \ : \ \mathcal{D}'_{\mathcal{O}}(K) \longrightarrow \mathcal{D}'_{\mathcal{O}}(K) \tag{19}$$

was defined in [9, § 5.4] and in [10, § 8]. (This Fourier transform is obtained as an application of double inductive limits from formula (10) to one-dimensional (usual) Fourier transform applied to the spaces $\mathcal{D}'(\mathcal{O}(s)/\mathcal{O}(r))$ for all pairs of integers $r < s$.) Note that for the two-dimensional Fourier transform from formula (19) it is important the self-duality of the additive group of the field K which is described above.

From [9, Prop. 25] and [10, Prop. 16] we obtain (after restriction to the space $\mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}$) that the map $\mathbf{F}$ is well-defined as a map

$$\mathbf{F} \,:\, \mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*} \longrightarrow \mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}.$$

In what follows we will consider only the last map which we denote by the same letter $\mathbf{F}$ as original map (19). We obtain the following properties from [9, Prop. 24, 25] and [10, Prop. 15, 16]:

$$\mathbf{F}^2 = \mathrm{id} \qquad \text{and} \qquad (a,b,c)\circ\mathbf{F} = \mathbf{F}\circ(-a,-b,c),$$

where $(a,b,c)\in G$. We note that the map $(a,b,c)\mapsto(-a,-b,c)$ is a second order automorphism of the group G.

Let $\varphi\in\mathcal{D}'(\bar{K})^{\mathcal{O}_{\bar{K}^*}}$. For any integer l we have the certain element φ_l (see formula (11)). By direct calculations we have

$$\mathbf{F}(\varphi_l) = (\mathbf{F}_{\mu_0}(\varphi))_{-1-l}, \tag{20}$$

where $\mu_0\in\mu(\bar{K})$ was defined in Remark 3. Hence we obtain that $\mathbf{F}(\Psi)=\Psi$.

Proposition 8. *We have the following properties.*

1. $\mathbf{F}(\beta(\Psi_1)) = \mathbf{F}(\beta(\Psi_q)) = \beta(\psi_1) = \beta(\psi_q)$.
2. $\mathbf{F}(\beta(\Psi_\alpha)) = \beta(\Psi_{q\alpha^{-1}})$ *for any* $\alpha\in\mathbb{C}^*$.

Proof. It follows from formula (20) and Proposition 7. The proposition is proved. □

Remark 4. Using Remark 3 and formula (20) we have $\mathbf{F}((g_\alpha)_l) = (g_{q\alpha^{-1}})_{-1-l}$ for $l\in\mathbb{Z}$ and $\alpha\in\mathbb{C}^*$. Inside the space $\beta(\Psi_1)$ we have $\mathbf{F}((\delta_0)_k) = (\delta_0)_{-k-2}$ for an integer k. Using that $(\delta_0)_k = (\mu_0)_{k+1}$ we have $\mathbf{F}((\mu_0)_k) = (\mu_0)_{-k}$.

5 Traces and "loop rotations"

We recall that in the proof of item 5 of Theorem 2 we have constructed a distinguished basis of the G-representation space Ψ_α (where $\alpha\in\mathbb{C}^*$) such that the basis consists of common eigenvectors for all elements h of the subgroup H and if $h\neq 0$ then distinct elements of the basis correspond to distinct eigenvalues of h. Note that this property defines the elements of this basis up to the multiplication on elements of $\mathbb{C}^*$. Elements $\Delta(l,g_\alpha)$ (where $l\in\mathbb{Z}$) which we considered above form this basis. Therefore the traces (when they exist) of elements of G with respect to the basis given by elements $\Delta(l,g_\alpha)$ (where $l\in\mathbb{Z}$) are invariants of G-representation Ψ_α.

Let $(a,b,c)\in G$. From formula (17) we obtain the following expressions for traces Tr_α in the G-representation Ψ_α:

- $\mathrm{Tr}_\alpha((a,b,c)) = 0$ if and only if $b\neq 0$;
- $\mathrm{Tr}_\alpha(a,0,c) = q^{-c-a}\alpha^a\sum\limits_{l\in\mathbb{Z}} q^{-al}$.

Clearly, the last expression is a divergent series.

To obtain convergent series as traces we will extend the group G to the group $\tilde{G} = G \rtimes \mathbb{Z}$ which will naturally acts on the space Ψ as well. Moreover, the subspace $\beta(\Psi_\alpha) \subset \Psi$ for any $\alpha \in \mathbb{C}^*$ defines a subrepresentation of the group $\tilde{G}$. Hence we will obtain an action of the group $\tilde{G}$ on the isomorphic space Ψ_α (recall Theorem 2). To construct the group $\tilde{G}$ we will consider automorphisms of the central extension $\widetilde{K^*}$ that are analogs of automorphisms of "loop rotation" from the theory of loop groups. We note that an extended discrete Heisenberg group was also considered in [11] to obtain the traces which are convergent series. But the methods of [11] were other than consideration of analogs of "loop rotations" and action on the space Ψ, which is a subspace of the space of distributions on a two-dimensional local field.

In what follows we will consider only a two-dimensional local field K that is isomorphic either to the first or to the second field from formula (4). Let t, u be a system of local parameters of K as it is described in the beginning of Section 3.1. For any $d \in \mathbb{Z}$ we consider an automorphism R_d of the field K given as:

- $R_d\big(\sum\limits_{i,j} a_{i,j} u^i t^j\big) = \sum\limits_{i,j} a_{i,j} u^{i+jd} t^j$, where $a_{i,j} \in \mathbb{F}_q$ and $K \simeq \mathbb{F}_q((u))((t))$;
- $R_d\big(\sum\limits_{j} b_j t^j\big) = \sum\limits_{j} b_j u^{jd} t^j$, where $b_j \in L$ and $K \simeq L((t))$ with L which is a finite extension of $\mathbb{Q}_p$.

It is clear that we have a homomorphism $\mathbb{Z} \to \mathrm{Aut}(K)$ given as $d \mapsto R_d$.

We note that $R_d(\mathcal{O}(i)) = \mathcal{O}(i)$ for any integer i. Hence we have a well-defined continuous automorphism R_d of the locally compact group $\mathcal{O}(i)/\mathcal{O}(j)$ for any integers $i \leq j$. This automorphism is the identity morphism on the group $\mathcal{O}(0)/\mathcal{O}(1)$. Thus, we obtain an action of the group $\mathbb{Z}$ on the $\mathbb{C}$-vector spaces $\mathcal{D}(\mathcal{O}(i)/\mathcal{O}(j))$ and $\mathcal{D}'(\mathcal{O}(i))/\mathcal{O}(j)$ via the maps induced by the maps R_d (which we will denote by the same letters R_d).

Since the automorphism R_d maps an element from $\mathcal{D}'(\mathcal{O}(i))/\mathcal{O}(j)$ that is invariant under all translations from $\mathcal{O}(i)/\mathcal{O}(j)$ to an element that is invariant under all translations from $\mathcal{O}(i)/\mathcal{O}(j)$, we obtain that R_d induces an automorphism of the one-dimensional $\mathbb{C}$-vector space $\mu(\mathcal{O}(i)/\mathcal{O}(j)) = \mathcal{D}'(\mathcal{O}(i)/\mathcal{O}(j))^{\mathcal{O}(i)/\mathcal{O}(j)}$. Therefore we have a well-defined action of the group $\mathbb{Z}$ on one-dimensional $\mathbb{C}$-vector spaces $\mu(\mathcal{O}(r) \mid \mathcal{O}(s))$ (for any integers r and s) and on the $\mathbb{C}$-vector spaces $\mathcal{D}_{\mathcal{O}}(K)$ and $\mathcal{D}'_{\mathcal{O}}(K)$ via the maps induced by the maps R_d (which we denote by the same letters).

Lemma 2. *For any integers b and d we have*

$$R_d(\mu_{B,0,b}) = q^{-\frac{b(b-1)d}{2}} \mu_{B,0,b}.$$

Proof. We consider the case $b \geq 0$. The case $b < 0$ follows from analogous computations. We have

$$\begin{aligned} R_d(\mu_{B,0,1} \otimes \ldots \otimes \mu_{B,b-1,b}) &= R_d(\mu_{B,0,1}) \otimes \ldots \otimes R_d(\mu_{B,b-1,b}) = \\ &= \mu_{B,0,1} \otimes q^{-d} \mu_{B,1,2} \otimes \ldots \otimes q^{-d(b-2)} \mu_{B,b-2,b-1} \otimes q^{-d(b-1)} \mu_{B,b-1,b} = \\ &= q^{-\frac{b(b-1)d}{2}} \mu_{B,0,1} \otimes \ldots \otimes \mu_{B,b-1,b} = q^{-\frac{b(b-1)d}{2}} \mu_{B,0,b}. \end{aligned}$$

The lemma is proved. $\square$

From the construction of the central extension $\widehat{K^*}$ we have that the group $\mathbb{Z}$ acts on $\widehat{K^*}$ via the group automorphisms induced by the maps R_d (which we denote by the same letters). From Lemma 2 we have that this action is also an action on the central extension $\widetilde{K^*}$. Clearly, $R_d(\mathcal{O}'^*) = \mathcal{O}'^*$ for any $d \in \mathbb{Z}$. Hence the group $\mathbb{Z}$ acts on the group $G \simeq \widetilde{K^*}/\mathcal{O}'^*$ via the group automorphisms induced by the maps R_d (which we denote by the same letters). Using an explicit description of the isomorphism $G \simeq \widetilde{K^*}/\mathcal{O}'^*$ given in the proof of Proposition 3 we calculate now this action, i.e. the homomorphism $\mathbb{Z} \to \mathrm{Aut}(G)$. We have (using also Lemma 2):

$$R_d((a,b,c)) = R_d\left((u^a t^b, q^{-c}\mu_{B,0,b})\right) = \left(R_d(u^a t^b), q^{-c} R_d(\mu_{B,0,b})\right) =$$
$$= \left(u^{a+db} t^b, q^{-c-\frac{b(b-1)d}{2}}\mu_{B,0,b})\right) = \left(a + db, b, c + \frac{b(b-1)d}{2}\right). \quad (21)$$

Remark 5. We can consider also formula (21) as a formula which gives a homomorphism from the subgroup $\left\{\begin{pmatrix} 1 & d \\ 0 & 1 \end{pmatrix}\right\}_{d\in\mathbb{Z}}$ of $GL(2,\mathbb{Z})$ to the automorphism group $\mathrm{Aut}(G)$. The first named author proved in [8] that this homomorphism can be extended to an injective homomorphism $GL(2,\mathbb{Z}) \hookrightarrow \mathrm{Aut}(G)$. By means of the last homomorphism it was also proved in [8] that the group $\mathrm{Aut}(G)$ is isomorphic to $(\mathbb{Z}\oplus\mathbb{Z}) \rtimes GL(2,\mathbb{Z})$. We note that other isomorphism $\mathrm{Aut}(G) \simeq (\mathbb{Z}\oplus\mathbb{Z}) \rtimes GL(2,\mathbb{Z})$ was constructed earlier by P. Kahn in [5] by another methods.

Now, using the homomorphism $\mathbb{Z} \to \mathrm{Aut}(G)$ given by formula (21), by the usual way we construct the group $\tilde{G}$ that is the semidirect product $G \rtimes \mathbb{Z}$. Explicitly, elements of $\tilde{G}$ are integer quadruples (a,b,c,d), where $(a,b,c) \in G$ and $d \in \mathbb{Z}$. From formula (21) we obtain the following explicit multiplication law in $\tilde{G}$:

$$(a_1,b_1,c_1,d_1)(a_2,b_2,c_2,d_2) = \big((a_1,b_1,c_1)\, R_{d_1}((a_2,b_2,c_2)), d_1+d_2\big) =$$
$$= \left((a_1,b_1,c_1)\left(a_2 + d_1 b_2, b_2, c_2 + \frac{b_2(b_2-1)d_1}{2}\right), d_1+d_2\right) =$$
$$= \left(a_1+a_2+d_1b_2, b_1+b_2, c_1+c_2+\frac{b_2(b_2-1)d_1}{2} + a_1 b_2, d_1+d_2\right).$$

Remark 6. From explicit multiplication law in $\tilde{G}$ we see that the group $\tilde{G}$ can be also considered as a group of the following matrices with integer entries:

$$\begin{pmatrix} 1 & d & a & c \\ 0 & 1 & b & \frac{b(b-1)}{2} \\ 0 & 0 & 1 & b \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

From this description it is evident that the group $\tilde{G}$ is nilpotent of class at most 3. We consider elements $x = (0,0,0,1)$, $y = (0,1,0,0)$ and $z = (0,1,0,0)$ of the group $\tilde{G}$. Then we have $[x,y] = (1,0,-1,0)$ and $[[x,y],z] = (0,0,1,0)$. Therefore the length of the lower central series of $\tilde{G}$ equals 3. Consequently, the group $\tilde{G}$ is nilpotent of class 3.

The group K^* acts on the group K by multiplication. Evidently, for any $d \in \mathbb{Z}$, $f \in K^*$, $h \in K$ we have $R_d(f \cdot h) = R_d(f) \cdot R_d(h)$. Hence and from construction of the actions, for any $g \in \widehat{K^*}$ and any $S \in \mathcal{D}_{\mathcal{O}}(K)$, any $T \in \mathcal{D}'_{\mathcal{O}}(K)$ we have

$$R_d(g \cdot S) = R_d(g) \cdot R_d(S), \qquad R_d(g \cdot T) = R_d(g) \cdot R_d(T). \tag{22}$$

Therefore we have that $R_d\left(\mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}\right) = \mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}$ for any $d \in \mathbb{Z}$. From formulas (22) we have also that $R_d(g \cdot (R_d^{-1}(T))) = R_d(g) \cdot T$. The last formula implies that for any $d \in \mathbb{Z}$, any $(a, b, c) \in G$ and any $V \in \mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}$ we have

$$R_d((a, b, c) \cdot (R_d^{-1}(V))) = R_d((a, b, c)) \cdot V. \tag{23}$$

From formula (23) we obtain that the following action of the group $\tilde{G}$ on the $\mathbb{C}$-vector space $\mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}$ is well-defined, i.e it is an action of the group:

$$(a, b, c, d) \cdot V = (a, b, c) \cdot (R_d(V)),$$

where $(a, b, c, d) \in \tilde{G}$ and $V \in \mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}$.

We recall that in formula (11) and Definition 1 we defined elements $\varphi_l \in \mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}$ and the $\mathbb{C}$-vector subspace Ψ of the space $\mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}$.

Proposition 9. *The space Ψ is a $\tilde{G}$-invariant subspace of the space $\mathcal{D}'_{\mathcal{O}}(K)^{\mathcal{O}'^*}$ with the following explicit action*

$$(a, b, c, d) \cdot \varphi_l = q^{-c-a(l+1)-\frac{dl(l+1)}{2}} \cdot ((a + dl)(\varphi))_{b+l},$$

where $(a, b, c, d) \in \tilde{G}$ and $(a + dl)(\varphi) \in \mathcal{D}'(\bar{K})^{\mathcal{O}_{\bar{K}}^*}$ is an application of the action of the element $u^{a+dl} \in \bar{K}^*$ to the element $\varphi \in \mathcal{D}'(\bar{K})^{\mathcal{O}_{\bar{K}}^*}$.

Proof. By the definition of the action, direct computation and Proposition 5 we have

$$\begin{aligned}(a, b, c, d) \cdot \varphi_l = (a, b, c) \cdot (R_d((0, l, 0) \cdot \varphi_0)) = (a, b, c) \cdot (R_d((0, l, 0)) \cdot R_d(\varphi_0)) = \\ = \left((a, b, c)\left(dl, l, \frac{dl(l-1)}{2}\right)\right) \cdot \varphi_0 = \left(a + dl, b + l, c + \frac{dl(l-1)}{2} + al\right) \cdot \varphi_0 = \\ = q^{-c-a(l+1)-\frac{dl(l+1)}{2}} \cdot ((a + dl)(\varphi))_{b+l}.\end{aligned}$$

The proposition is proved. □

From this proposition for any $\alpha \in C^*$ we obtain that

$$(a, b, c, d) \cdot (g_\alpha)_l = q^{-c-a(l+1)-\frac{dl(l+1)}{2}} \alpha^{a+dl} (g_\alpha)_{l+b},$$

where elements g_α were constructed in Remark 1. Hence, by direct computation from the last formula we obtain the following theorem.

Theorem 3. *For any $\alpha \in \mathbb{C}^*$ the subspace $\beta(\Psi_\alpha)$ of the space Ψ is $\tilde{G}$-invariant. We have the traces Tr_α of elements of $\tilde{G}$ in $\beta(\Psi_\alpha)$ with respect to a distinguished basis $(g_\alpha)_l = \beta(\Delta(l, g_\alpha))$, where $l \in \mathbb{Z}$ (see discussion in the beginning of this section):*

- $\mathrm{Tr}_\alpha((a,b,c,d)) = 0$ *if and only if* $b \neq 0$ *;*
- $\mathrm{Tr}_\alpha(a,0,c,d) = q^{-c-a}\alpha^a \sum\limits_{l\in\mathbb{Z}} q^{-al-\frac{dl(l+1)}{2}}\alpha^{dl}$.

Remark 7. Clearly, the series for $\mathrm{Tr}_\alpha(a,0,c,d)$ is a convergent series when $d>0$. Moreover, $\mathrm{Tr}_\alpha(a,0,c,d) = q^{-c-a}\alpha^a \sum\limits_{l\in\mathbb{Z}} \tilde{q}^{l^2}\eta^l$, where $\eta = q^{-a}q^{-\frac{d}{2}}\alpha^d$, $\tilde{q} = q^{-\frac{d}{2}}$. The series $\sum\limits_{l\in\mathbb{Z}} \tilde{q}^{l^2}\eta^l$ is the classical Jacobi theta function (compare also with [11, Example]).

It is easy to see that $\mathrm{Tr}_{q\alpha}(a,o,c,d) = \alpha^d \mathrm{Tr}_\alpha(a,0,c,d)$. Since the traces are invariants of a $\tilde{G}$-representation $\beta(\Psi_\alpha)$ with respect to a distinguished basis, from Theorems 2 and 3 we obtain a proposition.

Proposition 10. *The family of* $\tilde{G}$*-representations* $\beta(\Psi_\alpha)$ *parametrized by elements* α *from the set* $\mathbb{C}^* \setminus \{1\} = \mathbb{P}^1(\mathbb{C}) \setminus \{0,1,\infty\}$ *is a family of pairwise non-isomorphic irreducible* $\tilde{G}$*-representations.*

References

[1] Arnal' S. A., Parshin A. N., *On irreducible representations of discrete Heisenberg groups*, Matematicheskie Zametki, 92 : 3 (2012), pp. 323-330; english translation in Mathematical Notes, 92:3 (2012), pp. 295-301.

[2] Beloshapka I. V., Gorchinskiy S. O. *Irreducible representations of finitely generated nilpotent groups*, e-print arXiv:math/1508.06808 [math.RT] (to appear in Sb. Math.).

[3] Bernstein J., *Draft of: representations of p-adic groups*, lectures at Harvard University written by K.Rumelhart (1992) available at http://www.math.tau.ac.il/ bernstei .

[4] Dixmier J., *Les* C^**-algèbres et leurs représentations*, Paris, Gauthier-Villars Éditeur, 1969.

[5] Kahn P., *Automorphisms of the discrete Heisenberg groups*, preprint, 2005, 7 pp., available at http://www.math.cornell.edu/m/People/Faculty/kahn

[6] Kapranov M., *Semiinfinite symmetric powers*, e-print arXiv:math/0107089 [math.QA].

[7] Osipov D. V., n*-dimensional local fields and adeles on* n*-dimensional schemes*, Surveys in Contemporary Mathematics, Edited by N. Young, Y. Choi; Cambridge University Press, London Mathematical Society Lecture Note Series, No. 347 (2007), pp. 131-164.

[8] Osipov D. V., *The discrete Heisenberg group and its automorphism group*, Matematicheskie Zametki, 2015, 98:1, pp. 152-155; english translation in Mathematical Notes, 2015, 98:1, pp. 185-188.

[9] Osipov D. V., Parshin A. N., *Harmonic analisys on local fields and adelic spaces. I*, Izvestiya RAN: Ser. Mat., 2008, 72:5, pp. 77-140; english translation in Izvestiya: Mathematics, 2008, 72:5, pp. 915-976.

[10] Osipov D. V., Parshin A. N., *Harmonic analisys on local fields and adelic spaces. II*, Izvestiya RAN: Ser. Mat., 2011, 75:4, pp. 91-164; english translation in Izvestiya: Mathematics, 2011, 75:4, pp. 749-814.

[11] Parshin A. N., *On holomorphic representations of discrete Heisenberg groups*, Funktsional'nyi Analiz i Ego Prilozheniya, 44:2 (2010), pp.92-96; english translation in Functional Analysis and Its Applications , 44:2 (2010), pp. 156-159.

[12] Parshin A. N., *Representations of higher adelic groups and arithmetic*, Proceedings of the International Congress of Mathematicians (Hyderabad, India, 19-27 August 2010), Volume 1: Plenary lectures and ceremonies, World Scientific, 2010, pp. 362-392.

[13] Parshin A. N., *Notes on the Poisson formula.* With an appendix by Irina Rezvyakova. Algebra i Analiz 23 (2011), no. 5, pp. 1-54; translation in St. Petersburg Math. J. 23 (2012), no. 5, pp. 781-818.

[14] Weil A., *Fonctions zêta et distributions*, Séminaire Bourbaki, vol. 9, Années: 1965/1966, Exp. No. 312, Soc. Math. France, Paris 1995.

[15] Weil A., *Basic Number Theory*, 3rd ed., Grundlehren Math. Wiss., vol. 144, Springer-Verlag, New York–Berlin 1974.

D. V. Osipov

Steklov Mathematical Institute of Russsian Academy of Sciences, ul. Gubkina 8, Moscow, 119991 Russia

National University of Science and Technology "MISiS", Leninsky Prospekt 4, Moscow, 119049 Russia

E-mail: d_osipov@mi.ras.ru

A. N. Parshin

Steklov Mathematical Institute of Russsian Academy of Sciences, ul. Gubkina 8, Moscow, 119991 Russia

E-mail: parshin@mi.ras.ru